

Math Diversion Problem 250

P. Reany

December 21, 2024

In mathematics, you don't understand things.
You just get used to them.
— John von Neumann

The YouTube video is found at:

Source: <https://www.youtube.com/watch?v=Pj7ju4ebtY0>
Title: Can We Solve A Transcendental Equation
Presenter: SyberMath

1 The Problem

Given the relation

$$e^x + x + 1 = 0, \tag{1}$$

find the values of x .

2 The Preparation

I intend to use the Lambert W function, which goes as follows:

$$ze^z = B, \tag{2}$$

then

$$z = W(B), \tag{3}$$

where there are domain constraints on B that we won't go into here. Warning: This can be a complicated (multi-valued) function to deal with.

3 The Solution

Let's begin by rewriting the Given equation of the form

$$e^x = -x - 1. \tag{4}$$

Now, we introduce a new variable y , defined by

$$y = x + 1, \quad x = y - 1. \quad (5)$$

Then (1) becomes

$$e^{y-1} = -y, \quad (6)$$

or

$$e^{-1}e^y = -y, \quad (7)$$

And one more alteration:

$$e^{-1} = -ye^{-y}. \quad (8)$$

Next, we apply the Lambert W function:

$$W(e^{-1}) = -y. \quad (9)$$

And finally,

$$x = -W(e^{-1}) - 1. \quad (10)$$