

# Math Diversion Problem 67

P. Reany

September 28, 2024

## Abstract

Here we use the unipodal algebra to assist in solving the problem, which is given to us on YouTube. Although I'm referring to the series under the name 'olympiad', the problems are from diverse sources as olympiads, entrance exams, SATs, and the like.

It's not that I'm so smart; it's just that

I stay with problems longer.

—Albert Einstein

The YouTube video is found at:

Source: <https://www.youtube.com/shorts/I0hQS0opzfs>

Title: Challenging Math Olympiad Question

Presenter: Soul Institution

## 1 The Problem

Given the relation

$$\sqrt{x} + \sqrt{x+1} + \sqrt{x+2} = \sqrt{x+7}, \quad (1)$$

find the values of  $x$ .

## 2 The Prerequisites: The unipodal algebra

This algebra is formed as the extension of the complex numbers by the number  $u$ , where  $u^2 = 1$ , and  $u$  commutes with the complex numbers. The number  $u$  is said to be 'unipotent'. The set of numbers constructed this way are the unipodal numbers, a particular such number is called a unipode. The main conjugation operator on unipode  $a$  is the unegation operator, written  $a^-$ . It does not affect complex numbers, but it sends every  $u$  to its negative. Hence, if  $a = x + yu$ , where  $x, y$  are complex numbers, then  $a^- = x - yu$ . Unegation distributes over addition and multiplication.

The following are some properties that will come in handy:

$$u^2 = 1, \quad (2a)$$

$$u_{\pm} \equiv \frac{1}{2}(1 \pm u), \quad (2b)$$

$$u_{\pm}^2 = u_{\pm}, \quad (2c)$$

$$u = u_+ - u_-, \quad (2d)$$

$$u_+ u_- = 0, \quad (2e)$$

$$u_+ + u_- = 1, \quad (2f)$$

$$u u_+ = u_+, \quad (2g)$$

$$u u_- = -u_-, \quad (2h)$$

$$(u_{\pm})^- = u_{\mp}. \quad (2i)$$

You should prove (2c) – (2i). By the way, these two special unipodes  $u_{\pm}$  square to themselves. Such numbers in a ring are referred to as *idempotents*. In the unipodal numbers they have no inverses. The fact that the unipodal number system is not a field is of little concern to me. In fact, most unipodes have inverses, so long as they are not multiples of one of the idempotents. If one needs field elements, the scalars of the unipodal numbers comprise the field of complex numbers.

Two often-used results are, for complex numbers  $w, z$  (which are used to convert unipodes between the bases  $\{1, u\}$  and  $\{u_+, u_-\}$ ):

$$w + zu = (w + z)u_+ + (w - z)u_-, \quad (3a)$$

$$wu_+ + zu_- = \frac{1}{2}(w + z) + \frac{1}{2}(w - z)u. \quad (3b)$$

Next, we learn how to take the ‘norm’ of a unipode. Let  $w$  be a unipode in standard basis, given by

$$w = a + bu, \quad (4)$$

where  $a, b$  are complex numbers. The ‘norm’ of  $w$  is given as<sup>1</sup>

$$ww^- = (a + bu)(a - bu) = a^2 - b^2. \quad (5)$$

Now, let  $y$  be a unipode in idempotent basis, given as

$$y = Au_+ + Bu_-, \quad (6)$$

where  $A, B$  are complex numbers. The ‘norm’ of  $y$  is given as

$$\begin{aligned} yy^- &= (Au_+ + Bu_-)(Au_- + Bu_+) = ABu_+ + ABu_- \\ &= AB(u_+ + u_-) = AB. \end{aligned} \quad (7)$$

---

<sup>1</sup>Calling  $ww^-$  a ‘norm’ is rather imprecise. In accordance with terminology used by G. Sobczyk, I will call  $|ww^-|^{1/2}$  the (unipodal) modulus, and  $ww^-$  the (unipodal) dimodulus of  $w$ . See the Appendix.

The unipodal algebra has two copies of the complex numbers, one for each component. In any true unipodal equation, the corresponding coefficients across the equal sign are equal to each other. This is similar to equating real and imaginary components across the equal sign in the complex algebra.

When I first used the unipodal algebra to solve polynomial equations (c. 1984-5), I used the Clifford 1 algebra over the complex numbers. The ‘1’ means one unit vector  $u$ . So, a Clifford 1 number  $c$  can be represented as

$$c = a + bu, \quad (8)$$

where  $a, b$  are complex numbers. Of course,  $u$  being a unit vector, then

$$u^2 = 1. \quad (9)$$

Now, the standard basis for this space is  $\{1, u\}$  and the scalars are the complex numbers. To extract the ‘scalar part’ of (8), we use the selection operator  $\langle \cdot \rangle$ , as follows:

$$\langle c \rangle = \langle a + bu \rangle = a, \quad (10)$$

One can also subscript the selector with a zero for the scalar part:

$$\langle c \rangle_0 = \langle a + bu \rangle_0 = a, \quad (11)$$

and with a ‘1’ for the vector part:

$$\langle c \rangle_1 = \langle a + bu \rangle_1 = bu. \quad (12)$$

When I adopted the name ‘unipodal algebra’ from a paper I cowrote with two other authors, I found a need to adopt new terminology for naming the scalar and vector parts. Just as complex numbers are composed of a real number times the unit ‘1’ and another real number times the unit imaginary  $i$ , the unipodal numbers are composed of a complex number times the unit ‘1’ and another complex number times the unipotent number  $u$ . The part of the unipode that does not contain the unipotent factor is called the ‘complex part’ of the unipode. The part that does contain the unipotent element factor is called the **uniplex part** of the unipode.

Now, before you complain that calling the scalar part of a unipode the ‘complex part’ is nonsense, I point out that in complex analysis, the nonimaginary part is referred to as the ‘real part’. Lastly, when I say the ‘uniplex part’ in this series of papers, I refer only to the coefficient of the non-scalar part, which is complex only. Thus the uniplex part of unipode  $c = a + bu$  is just  $b$ . Another way to think of the uniplex part of  $c$  is to take the scalar (or complex) part of  $cu$ .

$$\langle c \rangle_1 = \langle cu \rangle = \langle au + b \rangle = b. \quad (13)$$

Thus, one must be careful when I report I’m taking the uniplex part of a unipode (across all the papers I’ve written over the years), because at times it may contain that factor of  $u$  and at other times not. But like I said: In this series it will always mean only the scalar factor of the unipotent element.

Much of the algebraic power of the unipodal algebra comes from 1) it being able to switch the presentation of a unipode between the standard basis and the idempotent basis, the latter basis being well suited for taking powers and roots.

### 3 The Solution

The straightforward approach to this problem is to just start squaring until the result is a polynomial in  $x$  and then find the root of the polynomial. But here I'm committed to using the unipodal algebra to find the root or roots. What I have so far isn't elegant, but it does reveal some interesting features of the algebra.

This has been one of the most revealing problems on the nature of my using the unipodal algebra to solve these 'olympiad' problems. I proved a simple but important lemma. Additionally, I added some unipodal heuristics for consideration.

Now, it seems obvious to me that our 'first unipodes' should be

$$a = \sqrt{x} + \sqrt{x+1} u, \tag{14a}$$

$$b = \sqrt{x+7} - \sqrt{x+2} u. \tag{14b}$$

There lacks no abundance of ways to define these unipodes. I tried many and settled down on these.

Okay, here's my game plan:

- 1) We start off in the basement (square roots) and use the given information Eq. (1), which will then free us to go upstairs, where there are no more square roots of polynomials.
- 2) We look for some ways to relate the square roots or polynomials in  $x$  in terms of either scalars or at least pure unipodes.<sup>2</sup>
- 3) If we haven't found a way to solve the problem already at this point, we project the significant unipodes into the scalars (the complex numbers).<sup>3</sup>

To fulfil step 1), we should convert the unipodes to idempotent form:

$$a = (\sqrt{x} + \sqrt{x+1}) u_+ + (\sqrt{x} - \sqrt{x+1}) u_-, \tag{15a}$$

$$b = (\sqrt{x+7} - \sqrt{x+2}) u_+ + (\sqrt{x+7} + \sqrt{x+2}) u_-, \tag{15b}$$

---

<sup>2</sup>A pure unipode contains no unknowns of the problem.

<sup>3</sup>The answer you want may be a real number or maybe only an integer, but those are still technically complex numbers.

and then subtract them:

$$\begin{aligned} a - b &= (\sqrt{x} + \sqrt{x+1} - (\sqrt{x+7} - \sqrt{x+2})) u_+ \\ &\quad + (\sqrt{x} - \sqrt{x+1} - (\sqrt{x+7} + \sqrt{x+2})) u_- \end{aligned} \quad (16a)$$

$$\begin{aligned} &= (\sqrt{x} + \sqrt{x+1} - \sqrt{x+7} + \sqrt{x+2}) u_+ \\ &\quad + (\sqrt{x} - \sqrt{x+1} - \sqrt{x+7} - \sqrt{x+2}) u_- . \end{aligned} \quad (16b)$$

But according to (1), the  $u_+$  component of (16b) is identically zero, leaving us with

$$a - b = (\sqrt{x} - \sqrt{x+1} - \sqrt{x+7} - \sqrt{x+2}) u_- . \quad (17)$$

Thus, we have completed step 1) by using the given information.

Now, there ought to be a number of ways of exploiting this towards a solution. One way is the following:

$$(a - b)(a - b)^- = 0 . \quad (18)$$

And so we've mapped a unipode  $(a - b)$  to a scalar.<sup>4</sup> Hopefully, we'll get useful information this time by taking the di-moduli of  $a$  and  $b$ :

$$aa^- = (\sqrt{x} + \sqrt{x+1})(\sqrt{x} - \sqrt{x+1}) = -1 , \quad (19a)$$

$$bb^- = (\sqrt{x+7} - \sqrt{x+2})(\sqrt{x+7} + \sqrt{x+2}) = 5 , \quad (19b)$$

and we have mapped two more unipodes to scalars.

What information can we get from (18) by expanding it? Let's see.

$$(a - b)(a - b)^- = (a - b)(a^- - b^-) \quad (20a)$$

$$= aa^- - ba^- - ab^- + bb^- \quad (20b)$$

$$= aa^- - 2\langle ba^- \rangle_0 + bb^- . \quad (20c)$$

From this last result and from (18), we can write:<sup>5</sup>

$$\langle ba^- \rangle_0 = \frac{1}{2}(aa^- + bb^-) , \quad (21)$$

which we can consider a lemma of the unipodal algebra. On applying the values from (19a) and (19b), we have that

$$\langle ba^- \rangle_0 = \frac{1}{2}(-1 + 5) = 2 . \quad (22)$$

---

<sup>4</sup>By the way, the reason this is true is because the di-modulus of a unipode  $c = c_+ u_+ + c_- u_-$  is given as  $cc^- = c_+ c_-$ , and in this case,  $c_+$  is identically zero. Of course, it's true that  $cc^- = 0$  if either  $c_+$  or  $c_-$  is identically zero.

<sup>5</sup>As a reminder, the notation ' $\langle \dots \rangle_0$ ' means to take the scalar component of the contained unipode, when it's in standard basis.

This above equation has actual information in it that might be useful, so let's expand it to find out.

$$\langle ba^- \rangle_0 = \langle [\sqrt{x+7} - \sqrt{x+2}u] \times [\sqrt{x} + \sqrt{x+1}u]^- \rangle_0 \quad (23a)$$

$$= \langle [\sqrt{x+7} - \sqrt{x+2}u] \times [\sqrt{x} - \sqrt{x+1}u] \rangle_0 \quad (23b)$$

$$= \langle (\sqrt{x+7}\sqrt{x} + \sqrt{x+2}\sqrt{x+1}) - (\sqrt{x+1}\sqrt{x+7} + \sqrt{x+2}\sqrt{x})u \rangle_0 \quad (23c)$$

$$= \sqrt{x+7}\sqrt{x} + \sqrt{x+2}\sqrt{x+1}, \quad (23d)$$

and on applying (22), we have that<sup>6</sup>

$$\sqrt{x+7}\sqrt{x} + \sqrt{x+2}\sqrt{x+1} = 2. \quad (24)$$

And perhaps we've reached the end of the line for the unipodal methods for this problem, and we should at this point just square (24) and simplify. Thus,

$$x(x+7) + 2\sqrt{x+7}\sqrt{x}\sqrt{x+2}\sqrt{x+1} + (x+2)(x+1) = 4, \quad (25)$$

or

$$\sqrt{x}\sqrt{x+1}\sqrt{x+2}\sqrt{x+7} = 2 - (x^2 + 5x + 1). \quad (26)$$

Once more unto the breach!<sup>7</sup> Square both sides and expand.

$$x^4 + 10x^3 + 23x^2 + 14x = x^4 + 10x^3 + 23x^2 - 10x + 1. \quad (27)$$

And this reduces to

$$14x = -10x + 1. \quad (28)$$

And this finalizes to

$$x = \frac{1}{24}. \quad (29)$$

Perhaps now we can see in Eq. (27) just why there's only one root. It's because it loses its quadratic and higher terms and is thus reduced to the 'linear' equation (28).

## 4 Conclusion

Let's take a moment to briefly take stock of the unipodal techniques we've used so far in this series that have been useful (and add in one or two that might be useful in the future):

- 1) Forming the 'first unipode' wisely.
- 2) Taking roots or powers, especially on unipodes in the idempotent basis.

---

<sup>6</sup>By the way, WolframAlpha.com confirms that this equation has the same root as does the original, which is 1/24.

<sup>7</sup>It's William Shakespeare! Coefficient of plagiarism = 0.

- 3) ‘Flipping’ between bases.
- 4) Extracting the complex and/or uniplex parts across an equation.
- 5) Taking the ‘magnitude square’ of a unipode. For example, if  $X = x_0 + x_1u$ ,  $XX^- = x_0^2 - x_1^2$ , which is, of course, just a complex number. If two unipodes are equal, their square magnitudes are equal, and you are free to calculate their square magnitudes from either basis.
- 6) Comparing square magnitudes this way:  $X^n(X^-)^n = (XX^-)^n$ .
- 7) If  $A$  and  $B$  are equal unipodes in standard form, then  $\frac{a_0}{a_1} = \frac{b_0}{b_1}$ , but if they are in idempotent form, then  $\frac{a_+}{a_-} = \frac{b_+}{b_-}$ .
- 8) A unipode can be written in standard or idempotent basis, but there is nothing to stop us from expressing a unipode  $A$  as a combination of all four basis elements, like this:

$$A = A_1u_+ + A_2u_- + A_3u + A_4, \quad (30)$$

where the  $A$ ’s are complex numbers. The unipode  $A$  is perfectly acceptable. Typically, we aren’t even looking for a unipode as the end result, but one of its components. For example, let say that we wanted the  $A_+$  component of  $A$ . There’s a short way to get it and a long way to get it. The short way is to just multiply through by  $u_+$ :

$$A_+u_+ = [A_1u_+ + A_2u_- + A_3u + A_4]u_+, \quad (31)$$

which would give us

$$A_+ = A_1 + A_3 + A_4, \quad (32)$$

where I scraped off the idempotent factor. By the way, the  $A_-$  component would look like this:

$$A_- = A_2 - A_3 + A_4. \quad (33)$$

This is what the long way looks like:

$$A = A_1u_+ + A_2u_- + A_3(u_+ - u_-) + A_4(u_+ + u_-) \quad (34a)$$

$$= (A_1 + A_3 + A_4)u_+ + (A_2 - A_3 + A_4)u_-. \quad (34b)$$

Furthermore, we can add to these tricks all the techniques of real and complex number theory and ring theory.

## 5 Appendix

Here I want to present a bit more theory on the unipodal algebra. I’ve found the need to do this because the broader the space of algebra problems I try to

solve with the unipodal algebra, the broader the unipodal theory I find I need to draw upon. And if I have to, the reader has to as well.

**Note:** I will be making references to Garret Sobczyk's book *New Foundations in Mathematics, The Geometric Concept of Number* [1], particularly in the sections he has on the unipodal and hyperbolic numbers.

Let's begin with the algebra of the **hyperbolic** extension of the real numbers. We start with the real numbers  $\mathbb{R}$  and extend them by the unipotent element  $u$ . This is denoted by  $\mathbb{R}[u]$ . Thus, a typical **hyperbolic number**  $h$  in standard form could be

$$h = x + yu, \quad (35)$$

where  $x, y$  are real numbers. Flipping this to idempotent form, we get

$$h = h_+ u_+ + h_- u_- . \quad (36)$$

For considerations due to symmetric  $2 \times 2$  matrices, Sobczyk calls the process of going from (35) to (36) the *spectral decomposition* of (35) ([1], p. 33). I suppose we could call this the 'spectral basis'. However, we will stick with calling it the 'idempotent basis'.

Let  $w$  be a general unipodal number for starters. Now, if  $w$  is neither zero nor a multiple of one of the idempotents, then it will have an inverse. The easiest way to find the inverse of  $w$  is to cast it first into the idempotent basis, like this:

$$w = w_+ u_+ + w_- u_- . \quad (37)$$

Then its inverse is

$$w^{-1} = w_+^{-1} u_+ + w_-^{-1} u_- = \frac{1}{w_+} u_+ + \frac{1}{w_-} u_- . \quad (38)$$

Clearly, this inverse exists because neither  $w_+$  nor  $w_-$  is zero, which we know to be the case because if either of them were zero, then  $w$  in (37) would reduce to being a multiple of one of the idempotents, which we have disallowed.

Next, comes the important issue of defining some sort of magnitude on the unipodal numbers, starting with the hyperbolic numbers. For hyperbolic number  $h$ , we can define the **hyperbolic modulus** by ([1], p. 25):

$$|h|_h \equiv \sqrt{|hh^-|}, \quad (39)$$

where, of course,  $hh^-$  is a real number.

Now, we can define something similar for unipodal numbers, such as for unipodal number  $w$ , we can define the **unipodal modulus** by:

$$|w|_u \equiv \sqrt{|ww^-|}, \quad (40)$$

where, of course,  $ww^-$  is a complex number, but  $|w|_u$  is a real number.



Now, I know why the real numbers play such an important role in the hyperbolic numbers, that being its close association with the hyperbolic plane and Lorentzian geometry. But it's been my experience in using the unipodal algebra to solve problems, that magnitudes of them represented by real numbers have not played much, if any, role (at least so far). Therefore, I propose to define a more useful notion of modulus for what I'm doing.

For unipodal number  $w$ , we can define the **unipodal di-modulus** by:

$$\text{mod}(w) = ww^{-}, \quad (41)$$

where, of course,  $ww^{-}$  is a complex number. The meaning of 'di-modulus' is this: The 'di' part refers to two aspects of the complex number  $ww^{-}$ , that being its magnitude and complex phase. And by not introducing squareroots, we refrain from burdening the algebra with unnecessary algebraic complications.

**Theorem:** If  $w$  is a unipode such that

$$ww^{-} = 1, \quad (42)$$

then

$$w^{-1} = w^{-}. \quad (43)$$

**Proof:**

Clearly,  $w$  is not zero, nor is it a multiple of an idempotent. Let's prove this by contradiction. Assume that

$$w = \alpha u_+, \quad (44)$$

where  $\alpha$  is a complex number. Then

$$ww^{-} = (\alpha u_+)(\alpha u_-) = \alpha^2 u_+ u_- = 0. \quad (45)$$

But  $ww^{-}$  cannot be both unity and zero at the same time, hence, a contradiction. Therefore  $w$  is not a multiple of  $u_+$ ; and by a similar argument, it is not a multiple of  $u_-$ .

Thus we know that  $w^{-1}$  exists. Therefore, multiplying across (42), we have that

$$w^{-1}(ww^{-}) = (w^{-1}w)w^{-} = w^{-1}. \quad (46)$$

And thus,

$$w^{-} = w^{-1}. \quad (47)$$

**Lemma:** If  $a, b$  are unipodes, then

$$a^{-}b^{-} = (ab)^{-}. \quad (48)$$

**Proof:** Hint: Set  $a = a_+u_+ + a_-u_-$  and  $b = b_+u_+ + b_-u_-$  and work it out.

---

**Theorem:** If  $w$  is a unipode then for positive integer  $n$

$$(w^-)^n = (w^n)^-. \quad (49)$$

**Proof:** (By induction) For  $n = 1$  there's nothing to show.

Multiply (49) through by  $w^-$ :

$$w^-(w^-)^n = w^-(w^n)^-. \quad (50)$$

The LHS becomes  $(w^-)^{n+1}$  by ordinary product-counting rules. The RHS becomes  $(w^{n+1})^-$  by the previous lemma. Therefore,

$$(w^-)^{n+1} = (w^{n+1})^-. \quad (51)$$

So, by assuming that the rule is true for case  $n$ , we were able to show that the rule also works for case  $n + 1$ . And we're done.

---

**Definition:** A nonzero unipode that has a vanishing di-modulus is said to be **isotropic**.

**Review:** Let  $c$  be a nonzero unipode. There are basically two ways to calculate its di-modulus. First, when nonzero  $c$  is represented in standard form,<sup>8</sup> as

$$c = c_0 + c_1u, \quad (52)$$

then its di-modulus is given as

$$cc^- = (c_0 + c_1u)(c_0 - c_1u) = c_0^2 - c_1^2. \quad (53)$$

So, then  $c$  is isotropic if  $c_0$  and  $c_1$  have the same squares. But if nonzero  $c$  is expressed in idempotent form,<sup>9</sup> then

$$c = c_+u_+ + c_-u_-, \quad (54)$$

and its di-modulus is given as

$$cc^- = (c_+u_+ + c_-u_-)(c_+u_- + c_-u_+) = c_+c_-(u_+ + u_-) = c_+c_-. \quad (55)$$

And thus  $c$  is isotropic if either  $c_+$  or  $c_-$  is zero.

---

<sup>8</sup>It cannot be that both  $c_0$  and  $c_1$  are zero.

<sup>9</sup>It cannot be that both  $c_+$  and  $c_-$  are zero.

**Lemma:**

Let  $a$  and  $b$  be distinct unipodes. Then, we can confirm by simple calculation that<sup>10</sup>

$$(a - b)(a - b)^- = (a - b)(a^- - b^-) \tag{56a}$$

$$= aa^- - ba^- - ab^- + bb^- \tag{56b}$$

$$= aa^- - 2\langle ba^- \rangle_0 + bb^- . \tag{56c}$$

Now, if unipode  $a - b$  is isotropic, then

$$aa^- - 2\langle ba^- \rangle_0 + bb^- = 0. \tag{57}$$

We can solve for any of these three terms, but I'll solve for the middle one:

$$\langle ba^- \rangle_0 = \frac{1}{2}(aa^- + bb^-), \tag{58}$$

which has proven itself to be useful on the unipodal battlefield.

So, what's the application of this lemma? In the first place, this lemma assists us in mapping unipodal information into scalar (i.e., the complex numbers) information, which is more likely to be useful in terms of the results we are after.

Now, the problem solver is the one who invents the two 'first unipodes'  $a$  and  $b$  in the first place. (Though in most problems I've solved with the aid of the unipodal algebra, I needed only one 'first unipode'.) And if the problem solver is clever enough, he or she might be able to tweak  $a - b$  into an isotropic unipode, thereby allowing the use (58). Of course, I'm not claiming that (58) will automatically be useful, but it may be worth trying.

## References

- [1] G. Sobczyk, *New Foundations in Mathematics, The Geometric Concept of Number*, Birkhauser/Springer, New York, 2013.
- [2] D. Hestenes, P. Reany, G. Sobczyk, 'Unipodal Algebra and Roots of Polynomials', *Advances in Applied Clifford Algebras*, Vol. **1**, No. 1, 51–64, 1991, Mexico City.
- [3] P. Reany, 'Unipodal Algebra and 2nd-Order Linear Differential Equations', *Advances in Applied Clifford Algebras*, Vol. **3**, No. 2, 121–126, 1993, Mexico City.

---

<sup>10</sup>As a reminder, the notation ' $\langle \dots \rangle_0$ ' means to take the scalar component of the contained unipode, when it's in standard basis.