

Trigonometry

P. Reany

February 3, 2025

You cannot read mathematics the way you read a
novel. If you zip through a page in less than an
hour, you are probably going too fast.

— Sheldon Axler
(from *Linear Algebra Done Right*)

1 Trigonometry Presentation

Trigonometry is big subject that I can only present a bare minimum on. I will, however, present trigonometry (meaning trigonometric functions) from the standpoint of both the real and complex numbers.¹

Think of this as a reference guide rather than a tutorial.

2 Basics of Trig Functions over the Reals

Mostly, I'm just going to present the common definitions and identities.

Let's begin with the most famous trig identity, the Pythagorean Identity on the unit circle involving the cosine and sine functions:

$$\cos^2 \theta + \sin^2 \theta = 1. \quad (1)$$

Next, some definitions (in order of presentation: secant, cosecant, tangent, cotangent):

$$\sec \theta = \frac{1}{\cos \theta}, \quad (2a)$$

$$\csc \theta = \frac{1}{\sin \theta}, \quad (2b)$$

$$\tan \theta = \frac{\sin}{\cos \theta}, \quad (2c)$$

$$\cot \theta = \frac{\sin}{\cos \theta}. \quad (2d)$$

(2e)

¹I'll probably go into more depth as time goes on.

Next, some common secondary identities:

$$1 + \tan^2 \theta = \sec^2 \theta, \quad (3a)$$

$$1 + \cot^2 \theta = \csc^2 \theta. \quad (3b)$$

Now, we look at some phase shifts:

$$\sin\left(\theta \pm \frac{\pi}{2}\right) = \pm \cos \theta, \quad (4a)$$

$$\cos\left(\theta \pm \frac{\pi}{2}\right) = \mp \sin \theta, \quad (4b)$$

$$\tan\left(\theta \pm \frac{\pi}{4}\right) = \frac{\cot \theta \pm 1}{1 \mp \tan \theta}, \quad (4c)$$

$$\sin(\theta + \pi) = -\sin \theta, \quad (4d)$$

$$\cos(\theta + \pi) = -\cos \theta, \quad (4e)$$

$$\tan\left(\theta + \frac{\pi}{2}\right) = -\cot \theta. \quad (4f)$$

Next, angle sums and differences:

$$\sin(\alpha \pm \beta) = \sin \alpha \cos \beta \pm \cos \alpha \sin \beta, \quad (5a)$$

$$\cos(\alpha \pm \beta) = \cos \alpha \cos \beta \mp \sin \alpha \sin \beta, \quad (5b)$$

$$\tan(\alpha \pm \beta) = \frac{\tan \alpha \pm \tan \beta}{1 \mp \tan \alpha \tan \beta}. \quad (5c)$$

Now we look at a few special cases of double angles:

$$\sin(2\theta) = 2 \sin \theta \cos \theta = \frac{2 \tan \theta}{1 + \tan^2 \theta}, \quad (6a)$$

$$\cos(2\theta) = \cos^2 \theta - \sin^2 \theta = \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta}, \quad (6b)$$

$$\tan(2\theta) = \frac{2 \tan \theta}{1 - \tan^2 \theta}. \quad (6c)$$

Next, we look at some half-angle formulas:

$$\sin\left(\frac{1}{2}\theta\right) = \operatorname{sgn}\left(\sin\left(\frac{1}{2}\theta\right)\right) \sqrt{\frac{1 - \cos \theta}{2}}, \quad (7a)$$

$$\cos\left(\frac{1}{2}\theta\right) = \operatorname{sgn}\left(\cos\left(\frac{1}{2}\theta\right)\right) \sqrt{\frac{1 + \cos \theta}{2}}, \quad (7b)$$

$$\tan\left(\frac{1}{2}\theta\right) = \frac{\sin \theta}{1 + \cos \theta} = \frac{1 - \cos \theta}{\sin \theta} = \frac{\tan \theta}{1 + \sec \theta}. \quad (7c)$$

3 Basics of Complex Numbers with Trig Functions

Let's begin with the Euler relations:

$$\cos \theta + i \sin \theta = e^{i\theta}, \quad (8a)$$

$$\cos \theta - i \sin \theta = e^{-i\theta}, \quad (8b)$$

Next, let's invert them:

$$\cos \theta = \frac{1}{2}(e^{i\theta} + e^{-i\theta}), \quad (9a)$$

$$\sin \theta = \frac{1}{2i}(e^{i\theta} - e^{-i\theta}), \quad (9b)$$

where, in the above cases, I used the usually understood real variable θ , but that can be replaced by the complex variable z . In fact, soon we will do so.

Okay, how to represent $\tan z$ by exponentials?

$$\tan z = \frac{\sin z}{\cos z} = \frac{1}{i} \frac{e^{iz} - e^{-iz}}{e^{iz} + e^{-iz}}. \quad (10)$$

Relation of the usual trig functions to the hyperbolic trig functions:

$$i \sin \theta = \sinh(i\theta), \quad (11a)$$

$$\cos \theta = \cosh(i\theta), \quad (11b)$$

$$i \tan \theta = \tanh(i\theta), \quad (11c)$$

$$-i \cot \theta = \coth(i\theta). \quad (11d)$$

4 Appendix: The Basics of Complex Numbers

Typically, we find a generic complex number denoted by the letter z , but one is free to choose other letters, as well. So, if z is a complex number, in general it has both real and imaginary parts:

$$z = a + bi, \quad (12)$$

where a, b are real components of basis vectors $1, i$. But they are also expressed as, respectively, the 'real' and 'imaginary' components of z .

Complex conjugation of complex number z is an operation that leaves real numbers alone but replaces the unit imaginary i with its negative, i.e., $-i$. The symbols most often used to represent complex conjugation are the $*$ and the overbar. I'll usually use the overbar. Thus, the complex conjugate of z in (12) is

$$\bar{z} = a - bi. \quad (13)$$

Obviously, the complex conjugation of a pure real number has no effect.

A funny thing happens when we multiply a complex number by its conjugate:

$$z\bar{z} = (a + bi)(a - bi) = a^2 + b^2. \quad (14)$$

So, $z\bar{z}$ is zero if and only if $z = 0$, otherwise, it's a positive real number.

Another funny thing happens when we add a complex number and its conjugate: we also get a real number. Let's see.

$$z + \bar{z} = (a + bi) + (a - bi) = 2a. \quad (15)$$

Why do we care about this? Because sometimes we need to map complex numbers into the real numbers to get information on the complex numbers. This problem will show you that.

I'm not going to prove this here, but every complex number can be expressed in exponential (or polar) form:

$$z = a + bi = \sqrt{a^2 + b^2}e^{i\theta} = (z\bar{z})^{1/2}e^{i\theta} = re^{i\theta}, \quad (16)$$

where we can think of r as the length of the complex numbers z or \bar{z} .

$$r \equiv (z\bar{z})^{1/2} \quad \text{or} \quad r^2 = z\bar{z} = |z|^2. \quad (17)$$

So, it will be good to know all this stuff in this section before you attempt to follow my solutions to these complex variables problems.

By the way, the complex numbers are what's called a *field*, so they can be added, subtracted, multiplied, and divided by each other (except you can't divide by zero, as usual). And, therefore, you can apply the quadratic formula to them! (Yay!)

Lemma 1: If a complex number z is equal to its own conjugate $z = \bar{z}$, it's real.

Lemma 2: If a complex number z is complex conjugated twice then there's no change: $\bar{\bar{z}} = z$.

Lemma 3: The complex conjugated of a product or a sum is the product or sum of the complex conjugates: $\overline{z_1 z_2} = \bar{z}_1 \bar{z}_2$ and $\overline{z_1 + z_2} = \bar{z}_1 + \bar{z}_2$.

Lemma 4: If $s, t \in \mathbb{R}$ and $z = s + ti$ then

$$i\bar{z} = t + si. \quad (18)$$